

Social and ecological drivers of success in agri-environment schemes: the roles of farmers and environmental context

Article

Accepted Version

McCracken, M. E., Woodcock, B. A., Lobley, M., Pywell, R. F., Saratsi, E. ORCID: <https://orcid.org/0000-0001-5917-6463>, Swetnam, R. D., Mortimer, S. R., Harris, S. J., Winter, M., Hinsley, S. and Bullock, J. M. (2015) Social and ecological drivers of success in agri-environment schemes: the roles of farmers and environmental context. *Journal of Applied Ecology*, 52 (3). pp. 696-705. ISSN 0021-8901 doi: <https://doi.org/10.1111/1365-2664.12412> Available at <https://centaur.reading.ac.uk/40882/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://www.journalofappliedecology.org>

To link to this article DOI: <http://dx.doi.org/10.1111/1365-2664.12412>

Publisher: Wiley-Blackwell

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Social and ecological drivers of success in agri-environment schemes: the roles of farmers and environmental context

Morag E. McCracken¹, Ben A. Woodcock¹, Matt Lobley², Richard F. Pywell¹, Eirini Saratsi², Ruth D. Swetnam^{1,3}, Simon R. Mortimer⁴, Stephanie J. Harris⁴, Michael Winter², Shelley Hinsley¹, and James M. Bullock^{1*}.

¹NERC Centre for Ecology and Hydrology, Benson Lane, Wallingford, UK, OX10 8BB

²Centre for Rural Policy Research, Department of Politics, University of Exeter, Rennes Drive, Exeter, UK, EX4 4RJ

³Department of Geography, Staffordshire University, Leek Rd, Stoke-on-Trent, UK, ST4 2DF

⁴School of Agriculture, Policy and Development, University of Reading, Earley Gate, Reading, UK, RG6 6AR

***Corresponding author:** James Bullock; NERC Centre for Ecology and Hydrology, Benson Lane, Wallingford, UK, OX10 8BB; jmbul@ceh.ac.uk; +44 (0)7824 460866

Running title: The success of agri-environment schemes

Word count: 7,007

Number of: Tables 2; Figures 2; References 40

Summary

1. Agri-environment schemes remain a controversial approach to reversing biodiversity losses, partly because the drivers of variation in outcomes are poorly understood. In particular, there is a lack of studies that consider both social and ecological factors.

2. We analysed variation across 48 farms in the quality and biodiversity outcomes of agri-environmental habitats designed to provide pollen and nectar for bumblebees and butterflies or winter seed for birds. We used interviews and ecological surveys to gather data on farmer experience and understanding of agri-environment schemes, and local and landscape environmental factors.

3. Multimodel inference indicated social factors had a strong impact on outcomes and that farmer experiential learning was a key process. The quality of the created habitat was affected positively by the farmer's previous experience in environmental management. The farmer's confidence in their ability to carry out the required management was negatively related to the provision of floral resources. Farmers with more wildlife-friendly motivations tended to produce more floral resources, but fewer seed resources.

4. Bird, bumblebee and butterfly biodiversity responses were strongly affected by the quantity of seed or floral resources. Shelter enhanced biodiversity directly, increased floral resources and decreased seed yield. Seasonal weather patterns had large effects on both measures. Surprisingly, larger species pools and amounts of semi-natural habitat in the surrounding landscape had negative effects on biodiversity, which may indicate use by fauna of alternative foraging resources.

5. *Synthesis and application.* This is the first study to show a direct role of farmer social variables on the success of agri-environment schemes in supporting farmland biodiversity. It suggests that farmers are not simply implementing agri-environment options, but are

learning and improving outcomes by doing so. Better engagement with farmers and working with farmers who have a history of environmental management may therefore enhance success. The importance of a number of environmental factors may explain why agri-environment outcomes are variable, and suggests some – such as the weather – cannot be controlled. Others, such as shelter, could be incorporated into agri-environment prescriptions. The role of landscape factors remains complex and currently eludes simple conclusions about large-scale targeting of schemes.

Keywords: birds; bumblebees; butterflies; experiential learning; farmer; farmland; habitat quality; interdisciplinary; landscape; multimodel inference

Introduction

Agri-environment schemes offer farmers financial incentives to adopt wildlife-friendly management practices, and are implemented in several parts of the world with the goal of reversing biodiversity losses (Baylis *et al.* 2008; Lindenmayer *et al.* 2012). These schemes are costly – the European Union budgeted €22.2bn for the period 2007–2013 (EU 2011) – and controversial. Controversy arises because researchers have reported variable success of agri-environment schemes in enhancing biodiversity (Kleijn *et al.* 2006; Batary *et al.* 2010b). It is clear that well-designed and well-managed options can benefit target taxa. For example, Pywell *et al.* (2012) found that options designed for birds, bees or plants had increased richness and abundance of both rare and common species. Baker *et al.* (2012) showed positive effects of options providing winter seed resources on granivorous bird populations. The question therefore arises – what causes variation in the success of agri-environment schemes?

Some options seem to work less well than others. Pywell *et al.* (2012) demonstrated that general compared to more targeted management had little effect in enhancing birds, bees and plants, while Baker *et al.* (2012) found that habitats providing breeding season resources for birds were less effective than those supplying winter food. But even within options there is great variation in biodiversity responses (Batary *et al.* 2010a; Scheper *et al.* 2013). There are several studies of the drivers of agri-environmental success (with success defined variously), but individual projects have looked at only one or a few drivers. In this paper we take a holistic approach by assessing a number of putative social and environmental constraints on success; specifically farmer experience and understanding,

landscape and local environment, and the weather. In doing so, we consider success in terms of both biodiversity outcomes and habitat quality.

Social scientists have long considered the role of the farmer in agri-environment schemes, but their questions have tended to focus on why farmers do or do not participate in the schemes (Wilson & Hart 2001; Wynne-Jones 2013) or how to change farmer behaviours in relation to environmental management (Burton & Schwarz 2013; de Snoo *et al.* 2013). There is a consensus that many farmers show limited engagement with the aims of agri-environment schemes (Wilson & Hart 2001; Burton, Kuczera & Schwarz 2008), leading to concern that this may jeopardize scheme success (de Snoo *et al.* 2013). There is, however, little direct evidence to link farmer understanding of, and engagement with, agri-environmental management with biodiversity outcomes on the farm (Lobley *et al.* 2013). Indeed, despite calls for more interdisciplinary social and ecological research into rural land use (Phillipson, Lowe & Bullock 2009) there is little such work in relation to agri-environment schemes.

Much ecological work has focused on the roles of landscape and local environments in determining biodiversity outcomes. Several studies have shown that the abundance and diversity of target species in agri-environment habitats is greater: a) in landscapes with higher target species richness or amount of (semi-)natural habitat; and/or b) where local habitat quality (e.g. food plant diversity) is greater (Carvell *et al.* 2011; Concepcion *et al.* 2012; Shackelford *et al.* 2013). While weather conditions are rarely considered, it is likely that weather during surveys will affect animal activity and the weather during the preceding seasons will affect local population sizes (Pollard & Moss 1995).

106

107 While most studies focus on success in terms of biodiversity outcomes, the farmer can only
108 directly affect the quality of the created habitat. It is therefore useful to consider success in
109 these terms as well. In this paper we derive measures of habitat quality related to the
110 foraging resources made available to the target biota. As well as impacts of the farmer's
111 activities, such quality measures may be affected by local abiotic factors such as soil type,
112 shading and seasonal weather (Myers, Hokschi & Mason 2012).

113

114 Putting these social and ecological factors together, we hypothesize that the richness and
115 abundance of target taxa using agri-environment habitats are increased where: the
116 landscape contains more target species and semi-natural habitat, the quality of the created
117 habitat is higher and when weather conditions during the season and the survey period are
118 more optimal for these taxa. We expect local habitat quality to be important and
119 hypothesize that this is in turn affected by the farmer's experience in, and understanding of,
120 agri-environmental management, as well as local abiotic environmental factors. We
121 consider these hypotheses for agri-environment options developed to provide resources for
122 key declining taxa of the farmed environment: pollen and nectar for bees and butterflies;
123 and winter food for granivorous farmland birds.

Materials and methods

STUDY SITES AND AGRI-ENVIRONMENT OPTIONS

We assessed the success of two options available to arable farmers under the English Entry Level agri-environment scheme (ELS), which involve sowing selected plant species in 6 m wide strips at field edges. The Nectar Flower Mixture option NFM ('EF4' under ELS; Natural England (2013)) uses a mixture of at least three nectar-rich plant species to support nectar-feeding insects, specifically bumblebees and butterflies. The Wild Bird Seed Mixture WBM ('EF2' under ELS) requires at least three small-seed bearing plant species to be sown, and is designed to provide food for farmland birds, especially during winter and early spring (see Appendix S1 in Supporting Information for more detail). We assessed NFM and WBM because they had specific success criteria, in terms of the taxa targeted (Pywell *et al.* 2012).

We selected 48 arable or mixed farms that had NFM or WBM strips sown between autumn 2005 and autumn 2006. To represent a range of English farming landscapes, 24 farms were in the east (Cambridgeshire and Lincolnshire), which is flat with large arable fields, and 24 in the south-west (Wiltshire, Dorset, Devon & Somerset), which is more hilly, with smaller fields and more mixed arable and grass farms. Half of the farms in each region had NFM options and half WBM. All farms had a minimum of two fields with the relevant ELS option. The farms were selected: a) first by Natural England – the statutory body that manages ELS – examining their GENESIS database for farms meeting the required geographic, date and ELS option criteria; and then b) by contacting farmers until sufficient had been found that were willing to take part.

FARMER INTERVIEWS

Semi-structured interviews were conducted in 2007 with all farmers. The interviews were designed to explore farmer attitudes towards, and history of, environmental management and their perceptions and understanding of the management requirements for NFM or WBM. Loble *et al.* (2013) analysed these interviews, and we used them to calculate three measures of farmer attitudes to, and engagement with, agri-environment schemes. “Experience” describes, on a four point scale, the farmer’s history of environmental management both formally as part of a scheme and informally: some had long-lasting and frequent engagement (4); others less frequent engagement (3); while some had limited experience, perhaps undertaking a single project (2); and some had no previous engagement (1).

“Concerns” represents farmer statements about their perceptions as to how easy it would be to meet the stipulations for creating and managing the habitat (e.g. establishing the plants, limiting herbicide use, cutting requirements). Responses to each requirement were scored 1 (very difficult) to 5 (easy), and a mean score across requirements was derived for each farmer. Finally, “Motivation” categorized the farmers in terms of their stated motivation for where they placed the strips on the farm, from more wildlife-focused to more utilitarian. The three categories were: 1) the best for wildlife, 2) to fit in with farming operations, or 3) simply to fulfill ELS requirements. Spearman rank correlations across the 48 farms indicated that these measures were independent of each other. We did not consider the influence of farmer demographic variables (e.g. age or education) as these have a complex relationship with environmental behaviours (Burton 2014).

ECOLOGICAL SURVEYS

Ecological surveys were carried out in 2007 and repeated in 2008. Three strips – or two if there were no more – were surveyed on each farm and parallel measures were made in a nearby ‘control’ cropped area at a field edge and of equivalent size, shape and aspect. A shelter score (0–8) was calculated, which represented the number of directions in which the strip was protected by hedges, etc (Dover 1996). We obtained data from national sources further describing the physical environment of each strip: the Agricultural Land Classification ALC, which grades land from 1–5 according to its agricultural quality; and the soil type, which we classified into light, medium or heavy soils (see Appendix S2).

For NFM strips we counted the number of flower units (i.e. a single flower, a multi-flowered stem or an umbel; Heard *et al.* (2007)) and identified these to species in five 1 m² quadrats at 10 m intervals along two parallel 50 m transects during July and again in August (for later emerging species). Bumblebees (as colour groups, e.g. Heard *et al.* (2007) – for brevity we refer to these as species) and butterflies (to species) were surveyed along these transects by recording those foraging within a 4 m band centred on the transect. Insect surveys were carried out between 10:30 h and 17:00 h during dry weather at temperatures >16 °C, and weather conditions – air temperature and wind speed (from 0=calm to 5=strong breeze) – were recorded.

For WBM strips, we estimated the seed resource by gathering all seeds from each sown species in three 1 m² quadrats at 10 m intervals along two parallel 50 m transects in September. Samples were stored at -20 °C in the dark until processing, at which time the seeds were separated from other plant material, dried at 80 °C for 24 hr and weighed. Bird use of the whole strip was monitored in November, January and February, during weather

conductive to bird activity (e.g. avoiding rain or high winds). Timed bird counts were made from a distance and then all birds were flushed (Hinsley *et al.* 2010).

LANDSCAPE AND SEASONAL WEATHER VARIABLES

To describe the landscape context of each farm, land cover was mapped in a 4 x 4 km square centred on each farm using Google Earth and the CEH Land Cover Map 2007. We used this single square size and a single landscape measure – the percentage cover of semi-natural habitats (grassland, woods, heaths, etc) – to avoid type 1 errors and highly correlated variables. This scale encompasses foraging distances of the target taxa (e.g. Osborne *et al.* 2008), although the exact scale used was probably unimportant as differences among farms in % semi-natural cover were very similar for 2 x 2 km and 4 x 4 km squares (correlation coefficient = 0.81). Species pools were estimated from national datasets of species lists mapped on a 10 x 10 km grid (Appendix S2). The grid square overlapping the centre point of each farm was interrogated for species lists of: butterflies for the period 2005–2009; granivorous birds during the winter for 2007–2011; and bumblebees from 2000–2010.

Daily weather data through 2007 and 2008 were obtained from the British Atmospheric Data Centre for the weather station closest to each farm. Daily maxima or minima were averaged across specific seasons (winter = December–February, etc) according to hypotheses about how weather would affect certain response variables (e.g. winter bird numbers would be affected by winter minimum temperatures).

STATISTICAL ANALYSES

We analysed the success of NFM and WBM habitats in terms of: a) biodiversity responses and b) habitat quality in terms of resources for the target taxa. For a), we considered the number and species richness of butterflies, bumblebees and granivorous birds. Number was the sum across the multiple surveys in a year, and species the total seen across the surveys. For b), we considered the number and species richness of flowers (mean across the quadrats and surveys) and seed weight (mean across quadrats). Determinants of success were analysed using general linear mixed models in R (R_Core_Development_Team 2008) using the 'glme' function of the lme4 package (Bates 2010). The nine response variables were tested against subsets of continuous and categorical explanatory variables ('fixed effects': Tables 1, 2), which were selected to reflect our hypotheses about the roles of farmer and environmental factors. Note that because we included 'region' as a separate factor, any effects of other variables do not reflect differences between the south-western and eastern regions.

In addition to these fixed effects, year was treated as a repeated measure by nesting it as a random effect within a subject factor describing the smallest sampling unit, i.e. the individual strip. To account for additional random effects, replicate strips were nested within farm, allowing analysis of factors at both the farm and the strip scale (Table 1). All data were counts and were modelled using a Poisson error term with a log link function, with the exception of seed weight, which was $\ln(n+1)$ transformed and modelled with normal errors. When used as explanatory variables, seed weight and flower numbers were $\ln(n+1)$ transformed. For the analysis of seed weight responses, four outlier values (>1000 mg) were removed to improve model fit and ALC was excluded as performance of the mixed models showed it to be strongly collinear with other explanatory variables. Because birds

were surveyed over the whole strip we considered strip area in preliminary analyses, but this was collinear with other factors and had low importance and so was excluded from the full analyses.

We used multimodel inference, which allowed us to consider competing models and moderately collinear variables (Burnham & Anderson 2002; Freckleton 2011). For each response variable, models representing all possible combinations of the fixed effects (excluding interactions), including a null model and a saturated global model, were created and the AIC difference (Δ_i) was calculated as:

$$\Delta_i = AIC_i - AIC_{\min},$$

where AIC_{\min} is the lowest value of any model, and AIC_i is the model-specific value. Following Burnham and Anderson (2002), models with $\Delta_i < 4$ were considered to form a set that best explained variation. For this subset of R models, Akaike weights (w_i) were derived:

$$w_i = \frac{\exp\left[-\frac{1}{2}\Delta_i\right]}{\sum_{r=1}^R \exp\left[-\frac{1}{2}\Delta_r\right]},$$

where w_i represents the probability that model i would be the best fitting if the data were collected again under identical conditions. The relative importance of individual variables can be calculated as the w_i of all models within the $\Delta_i < 4$ subset sums to 1. The importance of individual fixed effects was assessed by summing the w_i values of all models containing that explanatory variable within the subset using the ‘MuMIn’ package (Bartoń 2013). As many variables were modelled, we focused subsequently on the most frequently-included variables with an importance ≥ 0.4 (all included variables are given in Tables 1, 2). Parameter estimates were weighted by w_i and averaged across all models. Following Symonds and

Moussalli (2011) we calculated the marginal R^2 value for the global model to indicate goodness of fit.

Results

The ELS strips were successful in that they had more target species and resources than the paired control (crop) strips. Generalized linear mixed models using Poisson errors and pairing ELS and control strips showed the former had higher bumblebee numbers (mean *per* strip, *per* year 10.6 vs. 0.3; $F_{1,242} = 686$, $P < 0.001$) and species (2.0 vs. 0.1; $F_{1,242} = 91$, $P < 0.001$), butterfly numbers (6.1 vs. 0.6; $F_{1,242} = 346$, $P < 0.001$) and species (2.2 vs. 0.5; $F_{1,242} = 75$, $P < 0.001$), flower numbers (672 vs. 71; $F_{1,242} = 39676$, $P < 0.001$), granivorous bird numbers (63 vs. 1.7; $F_{1,230} = 2946$, $P < 0.001$) and species (4.4 vs. 1.1; $F_{1,230} = 150$, $P < 0.001$), and seed weight (124 vs. 0 g; $F_{1,230} = 2629$, $P < 0.001$).

BIODIVERSITY OUTCOMES

The agri-environment strips had a wide range of bumblebee numbers (*per* strip, *per* year; 0–97) and species (0–6), butterfly numbers (0–50) and species (0–8), and granivorous bird numbers (0–485) and species (0–13). The global models explained variation in each response quite well ($R^2 = 0.28$ – 0.68), and to a similar extent to other large-scale agro-ecology studies (Gabriel *et al.* 2010). The most important explanatory variables were those describing the local environment (Table 1). Bumblebees, butterflies and birds were more abundant and diverse in strips which had more abundant and diverse flowers or a greater seed mass (Fig. 1), and in strips which were more sheltered. Weather conditions during the survey had generally minor importance, which may be because the surveys were done during a narrow set of benign conditions. Unsurprisingly, farmer social variables had little

direct importance for biodiversity measures although there were more bumblebee numbers and species on farms with more experienced farmers, and more butterfly species where farmers placed their strips in locations they considered best for wildlife.

Region had contrasting effects, with south-western farms having more bumblebee numbers and species, fewer butterfly numbers and species, and similar bird numbers and species to eastern farms. Landscape factors were often important, in that both the percentage of semi-natural habitat and the size of the species pool had (surprisingly) negative relationships with biodiversity. Bird numbers and species were enhanced under higher winter minimum temperatures, and a similar pattern was seen for insect numbers in relation to summer maximum temperatures.

HABITAT QUALITY OUTCOMES

There was large variation among strips in flower number (*per strip, per year*; 0–9329) and species (0–17), or seed weight (0–597 mg). No model explained variation in flower species richness in the NFM strips well ($R^2 \leq 0.06$), and no variable had high importance (Table 2). Models for flower number and seed weight performed better. According to these, more experienced farmers produced strips with more resources (Fig. 2). Higher flower numbers were also found on strips created by farmers who placed them on the basis of wildlife-focused than utilitarian motives, but the opposite pattern was shown for seed weight. Interestingly, farmers who had envisioned greater problems with establishing and maintaining these habitats produced strips with a greater seed yield. Of the environmental factors, region had little importance and the local conditions were important only in determining flower numbers, which were greater on sites of poorer agricultural quality and

which were more sheltered. Flower numbers and seed weight were boosted by higher maximum temperatures in the season preceding maturation of flowers (spring) or seeds (summer). In addition, flower numbers were negatively affected by higher temperatures in the summer.

Discussion

As we hypothesized, the biodiversity outcomes of the agri-environment schemes were influenced by a range of factors, including landscape variables, the quality of the local habitat, seasonal weather and conditions during the surveys. Habitat quality itself – i.e. floral or seed resources – responded to the farmers’ experience and understanding of agri-environmental management as well as local environment and seasonal weather. Below we consider the factors in detail, but this study has highlighted the importance of multiple drivers in explaining variation in the success of agri-environment schemes. This builds on previous work, which has shown that a suite of factors are required for agri-environment success, including relevant prescriptions, adequate management and proximity to source populations (Whittingham 2011; Pywell *et al.* 2012). We have for the first time demonstrated the direct roles of social alongside these ecological factors. This interdisciplinary insight suggests actions to improve the success of agri-environment schemes need to consider farmers’ motivations, landscape factors and the local environment.

FARMER EXPERIENCE AND UNDERSTANDING

While social scientists have researched farmers’ attitudes and motivations towards agri-environmental management (de Snoo *et al.* 2013), little is known about whether and how

these social drivers affect biodiversity outcomes. The social and natural sciences have different research traditions, and while there are a number of studies which have used interdisciplinary approaches (Phillipson, Lowe & Bullock 2009; Austin, Raffaelli & White 2013) there is still little work linking social and ecological data in quantitative analyses. Interviews provide complex qualitative data, and those with our farmers revealed a range of previous engagement with agri-environmental management, a variety of opinions about the ease with which farmers felt they would be able to implement the required management, and different motivations for taking part (Lobley *et al.* 2013). The social scientists in the project team translated these qualitative responses into quantitative scores, which allowed us to combine social with ecological data in linear mixed models.

This approach proved to be powerful in linking biodiversity outcomes to farmer motivations. In the agri-environment options investigated, farmers are asked to establish specific seed mixes in field margins, which supply food resources to the target taxa. Farmers with greater agri-environmental experience produced strips with more of these resources. Experience was scored relative to the length of time and frequency with which farmers stated they had been involved in environmental management. Agri-environment schemes such as that in England, which simply pay farmers to follow specific prescriptions, have been criticized as not actively engaging farmers or allowing them to develop skills in environmental management (Burton, Kuczera & Schwarz 2008; de Snoo *et al.* 2013). In our case, it seems that farmers had developed such skills through their involvement in agri-environmental management.

The unexpected findings that more experienced farmers had more bumblebees and more wildlife-focused farmers had more butterfly species on their strips independent of their effects on habitat quality raises the tantalizing prospect that more continuous agri-environmental management had allowed populations to increase. While this interpretation is speculative, it reflects the scheme's aim to facilitate population recovery of target species (Baker *et al.* 2012).

The fact that farmers with more concerns about the ease of management produced greater quantities of seed suggests that if farmers are learning experientially (Riley 2008) then this is more successful if they are aware of their own knowledge gaps. That is, those who thought it would be easy had a misplaced confidence. The conflicting effects of farmer motivation for strip placement on the quality of the two strip types may reflect the relative levels of knowledge about these habitats. NFM was quite novel for many farmers and so those more motivated by wildlife benefits may have managed these strips more carefully. Farmers are more familiar with the requirements for WBM as many sow game cover, which is similar. While the differences were small, utilitarian farmers achieved better WBM results.

The three social variables were not correlated and so these relationships reveal different aspects of the agri-environmental role of farmers. We did not link these social variables to specific actions carried out by the farmer. This was because: a) we did not want to burden farmers with recording their actions or to influence their behaviours by doing so; and b) we were more interested in the farmers' experience and motivations than the well-studied issue of how management affects outcomes. However, it is clear that we are only beginning to understand the role of farmers in achieving agri-environmental success.

LOCAL AND LARGE-SCALE ENVIRONMENTAL FACTORS

The agri-environmental prescriptions were supported by the importance of the abundance and richness of flowers in attracting bumblebees and butterflies (Carvell *et al.* 2011) and of seed resources in attracting granivorous birds (Hinsley *et al.* 2010). Shelter benefits animals by providing warmth and protection (Pywell *et al.* 2004). Our findings of a positive effect of shelter on flower numbers, but a negative effect on seed weight are more novel, and may reflect a balance of competition (e.g. shading) and facilitation (e.g. warming). More flowers under conditions of low agricultural quality (i.e. low ALC) may reflect lower cover of competitive grasses, etc (Pywell *et al.* 2005).

Several studies have found that bee and bird abundance and richness are higher within agri-environmental options in landscapes with more semi-natural habitat (Concepcion *et al.* 2012; Shackelford *et al.* 2013). There is less information on the role of the species richness in the landscape, although Pywell *et al.* (2012) found this had a positive effect for bees but none for birds. By contrast, our study suggested negative effects of the proportion of semi-natural habitat and/or the size of the species pool on all but one of the biodiversity measures. Some studies have shown that agri-environmental options can have smaller effects on biodiversity in more diverse landscapes, presumably because these offer alternative foraging resources (Batary *et al.* 2010a; Carvell *et al.* 2011). In our case it may be that smaller species pools and areas of semi-natural habitat indicate fewer alternative resources and so the agri-environment strips act as ‘honey pots’ in attracting more birds or insects. Whatever the mechanism, landscape effects on agri-environmental outcomes are not straightforward.

409

410 Seasonal weather effects on abundance of the target biota and floral and seed resources are
411 not surprising and reflect fundamental biological optima (Anguilletta 2009). However, it is
412 important to note the importance of weather patterns for spatio-temporal variation in
413 success, and that these may cause apparent failures which are beyond anyone's control.

414

415 IMPLICATIONS FOR IMPROVING AGRI-ENVIRONMENT SUCCESS

416 Agri-environmental research needs to move on from the question that has predominated
417 for some time – 'do they work?' – to ask instead – 'what are the causes of variation in
418 success?'. While some factors that affect outcomes have been studied – such as landscape
419 context – this paper has shown that a holistic understanding of drivers is necessary. In
420 particular, we have demonstrated the role of the farmer. In implementing agri-environment
421 management, the farmer is not simply carrying out prescribed tasks, but is making decisions
422 which impact on success. The importance of experience suggests that farmers gain
423 experiential understanding of agri-environment management. This indicates scheme success
424 might be improved by ensuring farmers stay engaged and build up experience. Indeed,
425 Jarratt (2012) found that as farmers become more engaged in environmental-friendly
426 farming there is a willingness to take on more complex conservation activities. This leads to
427 the question whether actively training farmers in agri-environment management might
428 expedite such learning (Lobley *et al.* 2013). Indeed a review of the English scheme (Defra
429 2008) recommended that farmers should get increased advice, although it remains to be
430 seen whether this will be implemented.

431

The farmer has a role in choosing which agri-environment options to use, their placement on the farm and their establishment and management. Our study covered the latter two processes and these determined the quality of habitat produced and, ultimately, how many birds or insects used these strips. The fact that the amount of shelter affected both the quality and biodiversity outcomes suggests that farmers might be advised to consider this factor when deciding where to place strips. Similarly, pollen and nectar flower strips might be best placed on poorer quality land. Understanding of the role of the weather has a different implication, in that it can help farmers and others understand why agri-environment options may perform badly sometimes, much as crops do. Landscape factors have a complex role and the lack of general patterns (Batary *et al.* 2010a; Concepcion *et al.* 2012) suggests that any large-scale targeting of agri-environment schemes should be done with caution.

Acknowledgements

Many people helped with fieldwork, especially Sarah Hulmes, Lucy Hulmes, Rich Broughton Paul Bellamy, John Redhead and Jodey Peyton. Claire Carvell advised on surveys. The research was funded under the RELU programme (grant RES-227-25-0010). We thank the following for providing data: BTO, BWARS, Butterfly Conservation and the BADC. Natural England helped with finding farmers and we also thank the 48 farmers for their cooperation.

Data accessibility

- Social and environmental data: NERC-Environmental Information Data Centre
doi:10.5285/d774f98f-030d-45bb-8042-7729573a13b2 (McCracken *et al.* 2015)

References

- Angilletta, M.J. (2009) *Thermal adaptation: a theoretical and empirical synthesis*. Oxford University Press, Oxford.
- Austin, Z., Raffaelli, D.G. & White, P.C.L. (2013) Interactions between ecological and social drivers in determining and managing biodiversity impacts of deer. *Biological Conservation*, **158**, 214-222.
- Baker, D.J., Freeman, S.N., Grice, P.V. & Siriwardena, G.M. (2012) Landscape-scale responses of birds to agri-environment management: a test of the English Environmental Stewardship scheme. *Journal of Applied Ecology*, **49**, 871-882.
- Bartoń, K. (2013) MuMIn: multi-model inference, R package version 1.9.13.
- Batary, P., Andras, B., Kleijn, D. & Tscharntke, T. (2010a) Landscape-moderated biodiversity effects of agri-environmental management: a meta-analysis. *Proceedings of the Royal Society B-Biological Sciences*, **278**, 1894-1902.
- Batary, P., Baldi, A., Saropataki, M., Kohler, F., Verhulst, J., Knop, E., Herzog, F. & Kleijn, D. (2010b) Effect of conservation management on bees and insect-pollinated grassland plant communities in three European countries. *Agriculture Ecosystems & Environment*, **136**, 35-39.
- Bates, D.M. (2010) *lme4: Mixed-effects modeling with R*. Springer, London.
- Baylis, K., Peplow, S., Rausser, G. & Simon, L. (2008) Agri-environmental policies in the EU and United States: A comparison. *Ecological Economics*, **65**, 753-764.
- Burnham, K.P. & Anderson, D.R. (2002) *Model selection and multimodel inference: a practical information-theoretic approach*, 2nd edn. New York.
- Burton, R.J.F. (2014) The influence of farmer demographic characteristics on environmental behaviour: A review. *Journal of Environmental Management*, **135**, 19-26.

479 Burton, R.J.F., Kuczera, C. & Schwarz, G. (2008) Exploring farmers' cultural resistance to
 480 voluntary agri-environmental schemes. *Sociologia Ruralis*, **48**, 16-37.

481 Burton, R.J.F. & Schwarz, G. (2013) Result-oriented agri-environmental schemes in Europe
 482 and their potential for promoting behavioural change. *Land Use Policy*, **30**, 628-641.

483 Carvell, C., Osborne, J.L., Bourke, A.F.G., Freeman, S.N., Pywell, R.F. & Heard, M.S. (2011)
 484 Bumble bee species' responses to a targeted conservation measure depend on
 485 landscape context and habitat quality. *Ecological Applications*, **21**, 1760-1771.

486 Concepcion, E.D., Diaz, M., Kleijn, D., Baldi, A., Batary, P., Clough, Y., Gabriel, D., Herzog, F.,
 487 Holzschuh, A., Knop, E., Marshall, E.J.P., Tscharntke, T. & Verhulst, J. (2012)
 488 Interactive effects of landscape context constrain the effectiveness of local agri-
 489 environmental management. *Journal of Applied Ecology*, **49**, 695-705.

490 de Snoo, G.R., Herzon, I., Staats, H., Burton, R.J.F., Schindler, S., van Dijk, J., Lokhorst, A.M.,
 491 Bullock, J.M., Lobley, M., Wr6bka, T., Schwarz, G. & Musters, C.J.M. (2013) Toward
 492 effective nature conservation on farmland: making farmers matter. *Conservation*
 493 *Letters*, **6**, 66-72.

494 Defra (2008) Environmental Stewardship review of progress. Defra, London.

495 Dover, J.W. (1996) Factors affecting the distribution of satyrid butterflies on arable
 496 farmland. *Journal of Applied Ecology*, **33**, 723-734.

497 EU (2011) Is agri-environment support well designed and managed? European court of
 498 auditors special report no. 7/2011. European Commission, Luxembourg.

499 Freckleton, R. (2011) Dealing with collinearity in behavioural and ecological data: model
 500 averaging and the problems of measurement error. *Behavioral Ecology and*
 501 *Sociobiology*, **65**, 91-101.

502 Gabriel, D., Sait, S.M., Hodgson, J.A., Schmutz, U., Kunin, W.E. & Benton, T.G. (2010) Scale
 503 matters: the impact of organic farming on biodiversity at different spatial scales.
 504 *Ecology Letters*, **13**, 858-869.

505 Heard, M.S., Carvell, C., Carreck, N.L., Rothery, P., Osborne, J.L. & Bourke, A.F.G. (2007)
 506 Landscape context not patch size determines bumble-bee density on flower mixtures
 507 sown for agri-environment schemes. *Biology Letters*, **3**, 638-641.

508 Hinsley, S.A., Redhead, J.W., Bellamy, P.E., Broughton, R.K., Hill, R.A., Heard, M.S. & Pywell,
 509 R.F. (2010) Testing agri-environment delivery for farmland birds at the farm scale:
 510 the Hillesden experiment. *Ibis*, **152**, 500-514.

511 Jarratt, S. (2012) Linking the environmentally friendly farming careers of farmers to their
 512 effective delivery of wildlife habitats within the East of England. PhD, Nottingham.

513 Kleijn, D., Baquero, R.A., Clough, Y., Diaz, M., De Esteban, J., Fernandez, F., Gabriel, D.,
 514 Herzog, F., Holzschuh, A., Johl, R., Knop, E., Kruess, A., Marshall, E.J.P., Steffan-
 515 Dewenter, I., Tschardtke, T., Verhulst, J., West, T.M. & Yela, J.L. (2006) Mixed
 516 biodiversity benefits of agri-environment schemes in five European countries.
 517 *Ecology Letters*, **9**, 243-254.

518 Lindenmayer, D., Wood, J., Montague-Drake, R., Michael, D., Crane, M., Okada, S.,
 519 MacGregor, C. & Gibbons, P. (2012) Is biodiversity management effective? Cross-
 520 sectional relationships between management, bird response and vegetation
 521 attributes in an Australian agri-environment scheme. *Biological Conservation*, **152**,
 522 62-73.

523 Loble, M., Saratsi, E., Winter, M. & Bullock, J.M. (2013) Training farmers in agri-
 524 environmental management: the case of Environmental Stewardship in lowland
 525 England. *International Journal of Agricultural Management*, **3**, 12-20.

526 McCracken, M.E., Woodcock, B.A., Lobley, M., Pywell, R.F., Saratsi, E., Swetnam, R.D.,
 527 Mortimer, S.R., Harris, S.J., Winter, M., Hinsley, S., Bullock, J.M. (2015). Biodiversity,
 528 environmental and social data 2007-2008 for the project Improving the Success of
 529 Agri-environment Schemes (FarmCAT). NERC-Environmental Information Data
 530 Centre doi:10.5285/d774f98f-030d-45bb-8042-7729573a13b2

531 Myers, M.C., Hokschi, B.J. & Mason, J.T. (2012) Butterfly response to floral resources during
 532 early establishment at a heterogeneous prairie biomass production site in Iowa, USA.
 533 *Journal of Insect Conservation*, **16**, 457-472.

534 Natural England (2013) Entry Level Stewardship Handbook 4th edition.

535 Osborne, J.L., Martin, A.P., Carreck, N.L., Swain, J.L., Knight, M.E., Goulson, D., Hale, R.J. &
 536 Sanderson, R.A. (2008) Bumblebee flight distances in relation to the forage
 537 landscape. *Journal of Animal Ecology*, **77**, 406-415.

538 Phillipson, J., Lowe, P. & Bullock, J.M. (2009) Navigating the social sciences:
 539 interdisciplinarity and ecology. *Journal of Applied Ecology*, **46**, 261-264.

540 Pollard, E. & Moss, D. (1995) Historical records of the occurrence of butterflies in Britain:
 541 examples showing associations between annual number of records and weather.
 542 *Global Change Biology*, **1**, 107-113.

543 Pywell, R.F., Heard, M.S., Bradbury, R.B., Hinsley, S., Nowakowski, M., Walker, K.J. & Bullock,
 544 J.M. (2012) Wildlife-friendly farming benefits rare birds, bees and plants. *Biology*
 545 *Letters*, **8**, 772-775.

546 Pywell, R.F., Warman, E.A., Carvell, C., Sparks, T.H., Dicks, L.V., Bennett, D., Wright, A.,
 547 Critchley, C.N.R. & Sherwood, A. (2005) Providing foraging resources for bumblebees
 548 in intensively farmed landscapes. *Biological Conservation*, **121**, 479-494.

549 Pywell, R.F., Warman, E.A., Sparks, T.H., Grootorex-Davies, J.N., Walker, K.J., Meek, W.R.,
 550 Carvell, C., Petit, S. & Firbank, L.G. (2004) Assessing habitat quality for butterflies on
 551 intensively managed arable farmland. *Biological Conservation*, **118**, 313-325.

552 R_Core_Development_Team (2008) R: Version 2.12.2. A Language and Environment for
 553 Statistical Computing. R Foundation for Statistical Computing, Bristol, UK. URL
 554 <http://cran.r-project.org>.

555 Riley, M. (2008) Experts in their fields: farmer-expert knowledges and environmentally
 556 friendly farming practices. *Environment and Planning A*, **40**, 1277-1293.

557 Scheper, J., Holzschuh, A., Kuussaari, M., Potts, S.G., Rundlof, M., Smith, H.G. & Kleijn, D.
 558 (2013) Environmental factors driving the effectiveness of European agri-
 559 environmental measures in mitigating pollinator loss - a meta-analysis. *Ecology*
 560 *Letters*, **16**, 912-920.

561 Shackelford, G., Steward, P.R., Benton, T.G., Kunin, W.E., Potts, S.G., Biesmeijer, J.C. & Sait,
 562 S.M. (2013) Comparison of pollinators and natural enemies: a meta-analysis of
 563 landscape and local effects on abundance and richness in crops. *Biological Reviews*,
 564 **88**, 1002-1021.

565 Symonds, M.E. & Moussalli, A. (2011) A brief guide to model selection, multimodel inference
 566 and model averaging in behavioural ecology using Akaike's information criterion.
 567 *Behavioral Ecology and Sociobiology*, **65**, 13-21.

568 Whittingham, M.J. (2011) The future of agri-environment schemes: biodiversity gains and
 569 ecosystem service delivery? *Journal of Applied Ecology*, **48**, 509-513.

570 Wilson, G.A. & Hart, K. (2001) Farmer participation in agri-environmental schemes: towards
 571 conservation-oriented thinking? *Sociologia Ruralis*, **41**, 254-274.

572 Wynne-Jones, S. (2013) Ecosystem service delivery in Wales: evaluating farmers'
573 engagement and willingness to participate. *Journal of Environmental Policy &*
574 *Planning*, **15**, 493-511.

575

576 **Supporting Information**

577 Supporting information is supplied with the online version of this article.

578 **Appendix S1.** Description of Nectar Flower Mix and Wild Bird Mix options.

579 **Appendix S2.** Additional data sources.

580 Table 1. The importance of social and ecological drivers of biodiversity outcomes across 24 farms with agri-environment options targeted at
581 pollen and nectar feeding insects and 24 farms with options targeted at seed-eating birds. Importance was derived using Akaike weights (w_i)
582 following averaging of linear mixed models, and the parameter estimates (Param. est.) are weighted by w_i and averaged across models.
583 Categorical variables are marked * and the parameter estimates are given. The most important variables – with importance ≥ 0.4 – are
584 highlighted

| Level | Local (strip) environment | | | Landscape | | Farmer social | | | Seasonal weather | Weather during survey | | Region* E;SW |
|---|---------------------------|----------------|---------|------------------------|--------------|---------------|----------|----------------------|-------------------------|-----------------------|-------|-----------------|
| Variable | Flower # | Flower species | Shelter | % Semi-natural habitat | Species pool | Experience | Concerns | Motivation* 1;2;3 | Summer max. temperature | Temperature | Wind | |
| Response = Bumblebee numbers. Marginal R ² = 0.68. Models where Δ _{AIC} < 4 = 31 of 4096 | | | | | | | | | | | | |
| Importance | 1 | 1 | 0.55 | 1 | 0.63 | 1 | 0.18 | 0.13 | 0.32 | 1 | 0.40 | 0.83 |
| Param. est. | 0.46 | 0.11 | 0.11 | -0.06 | -0.07 | 0.28 | 0.13 | 0.06;0.32;0.46 | 0.21 | 0.17 | -0.07 | 0.06;0.49 |
| Response = Bumblebee species richness. Marginal R ² = 0.48. Models where Δ _{AIC} < 4 = 144 of 4096 | | | | | | | | | | | | |
| Importance | 1 | 0.57 | 0.95 | 0.31 | 0.53 | 0.63 | 0.12 | 0.05 | 0 | 0 | 0.15 | 0.74 |
| Param. est. | 0.25 | 0.04 | 0.1 | -0.01 | -0.03 | 0.13 | -0.02 | 0.38;0.41;0.15 | - | - | -0.04 | 0.38;0.56 |
| Response = Butterfly numbers. Marginal R ² = 0.28. Models where Δ _{AIC} < 4 = 99 of 4096 | | | | | | | | | | | | |
| Importance | 0.46 | 1 | 0.73 | 0.17 | 0.48 | 0.12 | 0.12 | 0.12 | 0.85 | 0.09 | 0.1 | 0.62 |
| Param. est. | 0.05 | 0.07 | 0.14 | -0.02 | -0.04 | -0.06 | 0.09 | 2.12;0.91;0.17 | 0.43 | 0.01 | -0.02 | 2.14;0.55 |
| Response = Butterfly species richness. Marginal R ² = 0.29. Models where Δ _{AIC} < 4 = 72 of 4096 | | | | | | | | | | | | |
| Importance | 0.71 | 0.15 | 0.83 | 0.27 | 0.18 | 0.38 | 0.08 | 0.87 | 0.12 | 0.08 | 0.08 | 1 |
| Param. est. | 0.07 | -0.02 | 0.09 | -0.01 | -0.02 | -0.12 | -0.01 | 1.44;0.53;0.69 | -0.09 | 0.01 | 0.1 | 1.43;0.13 |
| | | | | | | | | | | | | |
| Variable | Seed weight | | Shelter | % Semi-nat. habitat | Species pool | Experience | Concerns | Motivation* 1;2;3 | Winter min. temperature | N/A | | Region* E;SW |
| Response = Granivorous bird numbers. Marginal R ² = 0.36. Models where Δ _{AIC} < 4 = 19 of 512 | | | | | | | | | | | | |
| Importance | 1 | | 1 | 0.17 | 0.45 | 0.17 | 0.38 | 0.05 | 1 | | | 0.18 |
| Param. est. | 0.16 | | 0.27 | 0.01 | -0.14 | -0.09 | -0.23 | 8.8;12.5;14.5 | 0.46 | | | 8.8;7.5 |
| Response = Granivorous bird species richness. Marginal R ² = 0.36. Models where Δ _{AIC} < 4 = 41 of 512 | | | | | | | | | | | | |
| Importance | 1 | | 0.38 | 0.69 | 0.57 | 0.37 | 0.14 | 0.12 | 1 | | | 0.29 |
| Param. est. | 0.1 | | 0.04 | -0.02 | -0.06 | -0.08 | -0.04 | 0.92;1.03;1.18 | 0.21 | | | 0.92;0.79 |

Table 2. The importance of social and ecological drivers of habitat quality across 24 farms with agri-environment options targeted at pollen and nectar feeding insects (quality = flower numbers and species richness) and 24 with options targeted at seed-eating birds (quality = weight of seed). Importance was derived using Akaike weights (w_i) following averaging of linear mixed models, and the parameter estimates (Param. est.) are weighted by w_i and averaged across models. Categorical variables are marked with * and the parameter estimates are given. The most important variables – with importance ≥ 0.4 – are highlighted. ALC = Agricultural Land Classification

| Level | Local (strip) environment | | | Farmer social | | | Seasonal weather | | | Region* E;SW |
|---|---------------------------|--------------------------|---------|---------------|----------|----------------------|----------------------------|----------------------------|----------------------------|-----------------|
| Variable | ALC | Soil* Light;Med;Heavy | Shelter | Experience | Concerns | Motivation* 1;2;3 | Spring max. temperature | Summer max. temperature | | |
| Response = Flower numbers. Marginal R ² = 0.42. Models where Δ _{AIC} < 4 = 14 of 512 | | | | | | | | | | |
| Importance | 1 | 0.24 | 1 | 0.97 | 0.23 | 0.43 | 1 | 1 | | 0.34 |
| Param. est. | -0.72 | 963;720;1478 | 1.45 | 0.46 | 0.27 | 1477;720;166 | 4.5 | -0.31 | | 1477;741 |
| Response = Flower species richness. Marginal R ² = 0.06. Models where Δ _{AIC} < 4 = 49 of 512 | | | | | | | | | | |
| Importance | 0.35 | 0.04 | 0.22 | 0.36 | 0.21 | 0.01 | 0.21 | 0.21 | | 0.17 |
| Param. est. | 0.07 | 4.96;4.07;4.64 | -0.05 | 0.06 | 0.05 | 4.64;5.12;4.53 | 0.07 | 0.06 | | 4.64;4.81 |
| | | | | | | | | | | |
| Variable | ALC | Soil* Light;Med;Heavy | Shelter | Experience | Concerns | Motivation* 1;2;3 | Spring max. temperature | Summer max. temperature | Autumn max. temperature | Region* E;SW |
| Response = Seed weight. Marginal R ² = 0.21. Models where Δ _{AIC} < 4 = 55 of 512 | | | | | | | | | | |
| Importance | - | 0.01 | 0.43 | 0.7 | 0.74 | 0.64 | 0.15 | 0.71 | 0.31 | 0.19 |
| Param. est. | - | 167;166;191 | -11.5 | 36.4 | -35.1 | 191;200;299 | -6.45 | 46 | -35 | 191;184 |

Figure legends

Fig. 1. Examples of relationships between the major habitat quality drivers and biodiversity outcomes (see all drivers in Table 1). Circles show raw data, solid lines the fitted relationship (from linear mixed models, so accounting for other drivers) and dotted lines ± 1 standard error. a) Numbers of bumblebees, and b) Butterfly species richness as affected by the number of flowers. c) Numbers of seed-eating birds as affected by the weight of seeds. The unfilled circles in c) show large abundance values, which are, in order from left to right: 422, 485, 362, 223, 314 and 224.

Fig. 2. Examples of relationships between the length and intensity of the farmer's previous experience of environmental management (from 1 none to 4 high) and habitat quality measures in agri-environment strips (see all drivers in Table 2). Circles show raw data, solid lines the fitted relationship (from linear mixed models, so accounting for other drivers) and dotted lines ± 1 standard error of this fit. a) Number of flowers in a nectar flower strip. b) Weight of seeds in a wild bird seed strip. The unfilled circles show large values, which are: in a) 9329 and 5218; and in b) 597 mg.

Fig. 1




Fig. 2

